

ICD-10-CM Key Concepts to Consider when Documenting

In general, distinguish between primary, underlying, spontaneous, complication of, sequelae/late effect, manifestation of or secondary/due to.

<p>Type: <i>gestational</i> diabetes, <i>Greenstick</i> (closed) fracture, <i>neonatal</i> hypertension, <i>primary</i> osteoarthritis, <i>aspiration</i> pneumonia, <i>atypical</i> atrial fibrillation, <i>predominantly hyperactive</i> ADHD, <i>displaced comminuted</i> fracture, <i>cluster</i> headache, <i>benign</i> neoplasm of skin, <i>carcinoma in situ</i> exocervix, <i>systolic</i> congestive heart failure, <i>low tension</i> glaucoma, <i>periumbilic rebound</i> abdominal tenderness, <i>age-related</i> osteoporosis with current <i>pathological</i> fracture right humerus</p>	<p>Complications/Manifestations: hemorrhage, status asthmaticus, withdrawal, spontaneous rupture of eardrum, without goiter, Bouchard’s nodes, without aura, periodontal disease, with coma, gangrene, diabetic glaucoma, with intestinal obstruction</p>	<p>Associated Conditions/Cause/Contributing Factors: rheumatic, congenital, due to iodine deficiency, due to renal impairment, due to MRSA, due to insulin pump malfunction, due to streptococcus group B, due to excess calories, UTI associated with chronic Foley use, heart failure due to hypertension, age-related osteoporosis with current pathologic fracture, allergic reaction due to: cosmetics, plants, dyes, metals, adhesives, infectious agents</p> <p>It is important for the provider to document the association/correlation between the underlying condition and any associated conditions.</p>
<p>Severity: uncomplicated, mild persistent, intractable, with status migrainosus, mild stage, moderate stage, severe stage, indeterminate stage, stage 1, stage 5, first degree, 3rd degree</p>	<p>Anatomical Location: knee, great toe, thoracic spine, lower quadrant, upper eyelid, aortic valve, chewing surface, upper-inner quadrant, overlapping sites, anterior aspect, hard palate</p>	<p>Time Parameters: persistent, paroxysmal, acute, chronic, acute on chronic, recurrent, first trimester, gestational age 26 completed weeks, subacute, loss of consciousness for 30 minutes, no loss of consciousness</p>
<p>Episode: single, initial encounter, subsequent encounter, sequel/late effect</p>	<p>Localization: medial, proximal, central, peripheral, distal, lower end, etc.</p>	<p>Laterality: left, right, bilateral, generalized</p>
<p>Remission Status: full, partial</p>		